

Michigan
Pediatric Cardiac Protocols
PEDIATRIC WIDE COMPLEX TACHYCARDIA

Date: May 31, 2012

Page 1 of 2

Pediatric Wide Complex Tachycardia

Pre-Medical Control

PARAMEDIC

1. Follow the **Pediatric Assessment and Treatment Protocol**.

STABLE

1. Consider 12-Lead ECG, if available.

Post-Medical Control

2. Per MCA Selection Administer Lidocaine OR Amiodarone.

<u>Medication Options:</u> (choose one)	
<input type="checkbox"/>	Lidocaine 1 mg/kg IV/IO
OR	
<input checked="" type="checkbox"/>	Amiodarone 5 mg/kg IV/IO over 20-60 minutes


UNSTABLE

1. If Cardiopulmonary compromise is present as evidenced by hypotension, acutely altered mental status or signs of shock, contact medical control.

Post-Medical Control

2. If time and condition allow prior to cardioversion, sedate per MCA selection. Refer to **Patient Sedation Protocol**.
3. If HR greater than 180, consider Synchronized Cardioversion 0.5 – 1 J/kg.
4. Consider repeat cardioversions at 2 J/kg.
5. In borderline unstable patients, consider Adenosine if rhythm regular and QRS is monomorphic. If IV/IO is readily available, administer Adenosine 0.1 mg/kg (maximum 6 mg) IV/IO, rapid IV push through the most proximal injection site. This should be followed immediately with a 5 – 10 ml NS flush. May repeat Adenosine 0.2 mg/kg IV/IO (maximum 12 mg).

Michigan
Pediatric Cardiac Protocols
PEDIATRIC WIDE COMPLEX TACHYCARDIA



Medication Options:
(choose one)

Lidocaine 1 mg/kg IV/IO

OR

Amiodarone 5 mg/kg IV/IO over 20 – 60 minutes